

De Novo Acute Nonlymphoblastic Leukemia (M5b) Having a Chromosomal 3;21 Translocation With Immunoglobulin Heavy-Chain Gene Rearrangement

To the Editor: A reciprocal translocation between the long arms of chromosomes 3 and 21 with breakpoints at bands 3q26 and 21q22 [t(3;21)(q26;q22)] has been reported in malignant cells from patients with therapy-related myelodysplastic syndrome (t-MDS), therapy-related acute myeloid leukemia (t-AML), and chronic myelogenous leukemia (CML) [1-3], and it has been reported that the AML1-EVI-1 fusion gene is associated with this translocation [3,4]. However, we report on a rare patient who had de novo acute nonlymphoblastic leukemia with t(3;21) and another interesting feature.

A 67-year-old male was admitted in June 1995 with high fever. Annual medical examinations had revealed no hematologic disorders until then. Physical examination exhibited severe anemia. Hemoglobin was 58 g/l; platelets were $63 \times 10^9/l$; and white blood cell count was $77.2 \times 10^9/l$ with 82.5% blasts and 11% monocytes. The following blood chemistry values were recorded: lactate dehydrogenase, 1,230 IU/l; IgG, 3,467 mg/dl; IgA

1,050 mg/dl; IgM, 70 mg/dl; serum lysozyme, 64 μ g/ml. M protein, serum IL-1 β , and IL-6 were not detected.

Disseminated intravascular coagulation was not revealed. Bone-marrow aspirate was hypercellular, with 82.8% blasts which were negative in peroxidase stain and positive in alpha naphthyl butyrate esterase stain, and with 8% plasma cells which had mild dysplasia (Fig. 1). Surface antigen of blast cells was positive in CD13, CD14, CD33, CD34, CD19, and HLA-DR. Cytogenic studies performed on bone marrow showed 46XY,t(3;21) and Ig heavy-chain rearrangement, but major bcr by RT-PCR was not detected in peripheral blood cells. From the above findings, we diagnosed this case as de novo acute nonlymphoblastic leukemia M5b, with chromosomal abnormality 46XY,t(3;21) and B-lymphoid feature with mild plasma-cell dysplasia. This patient was treated with mitoxantrone (7 mg/m², days 1-3) and cytosine arabinoside (200 mg/m², days 1-5). He reached complete remission in October, 1995.

There were two notable points about this case. First, although the 3;21 translocation has been reported in patients with CML, t-AML, t-MDS, and MDS progressing to leukemia, it has very rarely been observed in de novo AML [3]. Secondly, the patient's bone-marrow aspiration revealed mild dysplasia of plasma cells. Akashi et al. [5] reported that simultaneous occurrence of myelomonocytic leukemia and multiple myeloma might originate from common leukemic progenitors. However, since the Ig heavy-chain gene rearrangement disappeared in the remission state of leukemia




Fig. 1. Bone-marrow aspirate showed leukemic cells without Auer bodies and plasma cells with mild dysplasia (Giemsa stain, $\times 1,000$). Nuclei of leukemic cells were just like those of monocytes, and had some nucleoli. There were giant plasma cells with multiple nuclei. M protein was never produced in these plasma cells.

in spite of remaining plasma-cell dysplasia, and since we failed to detect M protein during the patient's hospital course, these plasma cells were not regarded as a neoplasm. It was suggested that leukemic cells with CD19 surface antigen had an Ig heavy-chain gene rearrangement.

In summary, it is suggested that the 3;21 translocation might not rarely be detected also in de novo leukemia and might be associated with B-lymphoid features and plasma cell dysplasia.

HIROYASU KAYA
KEIJI AOSHIMA
KAZUHIRO OKAFUJI
NOBUYOSHI TANAKA

Department of Internal Medicine, Fukuiken Saiseikai Hospital,
Fukui, Japan

SHIGEKI OHTAKE
SHINOBU NAKAMURA
TAMOTSU MATSUDA

Third Department of Internal Medicine, Kanazawa University
School of Medicine, Ishikawa, Japan

REFERENCES

1. Rubin CM, Larson RA, Bitter MA, Carrino JJ, Le Beau MM, Diaz MO, Rowley JD: Association of a chromosomal 3;21 translocation with the blast phase of chronic myelogenous leukemia. *Blood* 70:1338-1342, 1987.
2. Rubin CM, Larson RA, Anastasi J, Winter JN, Thangavelu M, Vardiman JW, Rowley JD, Le Beau MM: t(3;21)(q26;q22): A recurring chromosomal abnormality in therapy-related myelodysplastic syndrome and acute myeloid leukemia. *Blood* 76:2594-2598, 1990.
3. Nucifora G, Rowley JD: AML1 and the 8;21 and 3;21 translocations in acute and chronic myeloid leukemia. *Blood* 86:1-14, 1995.
4. Mitani K, Ogawa S, Tanaka T, Miyoshi H, Kurokawa M, Mano H, Yazaki Y, Ohki M, Hirai H: Generation of the AML1-EVI-1 fusion gene in the t(3;21)(q26;q22) causes blastic crisis in chronic myelocytic leukemia. *EMBO J* 13:504-510, 1994.
5. Akashi K, Harada M, Shibuya T, Fukagawa K, Kimura N, Sagawa K, Yoshikai Y, Teshima T, Kikuchi M, Niho Y: Simultaneous occurrence of myelomonocytic leukemia and multiple myeloma: Involvement of common leukemic progenitors and their developmental abnormality of "lineage infidelity". *J Cell Physiol* 148:446-456, 1991.

Hematologic Benefits of 1-Hydroxyvitamin D₃ in an Elderly Patient With Chronic Myelodysplastic Syndrome

To the Editor: Myelodysplastic syndromes (MDS) represent a heterogeneous group of preleukemic disorders, characterized by advanced age-of-onset and peripheral cytopenias [1]. Among clinical treatments, it has been generally thought that 1,25-dihydroxyvitamin D₃ acts to induce differentiation of blast cells in MDS, and that a high dose of 1,25-dihydroxyvitamin D₃ produces hypercalcemia [2-4]. We present a patient with MDS who exhibited an elevation of a previously low platelet count and hemoglobin level after the administration of a low dose of 1-hydroxyvitamin D₃, such as used in treatment of osteoporosis.

A 70-year-old Japanese woman was admitted to our hospital with purpura on March 3, 1995. Platelet count was $1.0 \times 10^4/\mu\text{l}$, hemoglobin level was 8.7 g/dl, and mean corpuscular volume (MCV) was 107.7 fl. The reticulocyte count was $9.2 \times 10^4/\mu\text{l}$, and no nucleated red blood cells were found. Examination of a bone-marrow aspirate revealed normocellular marrow with degranulated granulocytes and pseudo-Pelger-Huët anomalies. Blasts constituted 2.4% of all nucleated cells. No chromosomal abnormalities were observed in 20 metaphase cells. Biochemical evaluation showed elevated serum levels of LDH (529 IU/l), hemoglobin-F (1.4%), ferritin (99 ng/ml), and folic acid (11.7 ng/ml), with a normal serum level of vitamin B12 (690 pg/ml). Indirect bilirubin was 0.8 mg/dl. Combs test was negative. The patient was seropositive for the surface antigen of the hepatitis B virus. The level of platelet-associated IgG (PAIgG) was markedly elevated ($138.6 \text{ ng}/10^7 \text{ cells}$). Refractory anemia was diagnosed according to the FAB classification. Because of the patient's advanced age and seropositivity for hepatitis B virus, she was administered only 1 μg of 1-hydroxyvitamin D₃ per day. The bicytopenia improved gradually without any side effects: the platelet count rose to $5.4 \times 10^4/\mu\text{l}$ and the hemoglobin to 11.1 g/dl, over 10 weeks, and the purpura disappeared. Serum levels of LDH and indirect bilirubin were reduced to 383 IU/l and 0.3 mg/dl, respectively. The PAIgG was reduced to 55.9 ng/ 10^7 cells . On continued treatment for 9 months, the platelet count was maintained at about $5.0 \times 10^4/\mu\text{l}$, and the hemoglobin at about 11 g/dl (Fig. 1). A bone-marrow aspirate performed




Fig. 1. Clinical course during treatment with 1-hydroxyvitamin D₃. Changes in LDH, indirect bilirubin, and platelet-associated IgG are shown at bottom.